Cosmological models evolving through the Big Bang

Martina Adamo

PhD advisors: Flavio Mercati, Ángel Ballesteros Castañeda

Quantum gravity phenomenology in the multi-messenger approach COST Action CA18108 – Second Training School Belgrade, 9/Sep/2022








A singularity can be regarded as a place where there is a breakdown of the classical concept of spacetime as a manifold with a pseudo-Reimannian metric. Because all known laws of physics are formulated on a classical spacetime background, they will all break down at a singularity. This is a great crisis for physics because it means that **one cannot predict the future**. One does not know what will come out of a singularity.

S. W. Hawking, "Breakdown of Predictability in Gravitational Collapse", Phys. Rev. D14, 246 (1976) Gravitational singularities are regions of spacetime where geometry or other fundamental physical structures become meaningless, and this happens in a **coordinate-independent way**

- the volume goes to zero
- some eigenvalues of the energy-momentum tensor diverge
- some curvature invariants diverge
- the geodesic equations are singular

But does this imply that dynamics is not well-defined? Is this enough to give up on classical determinism?

Classical determinism in General Relativity



<u>GR:</u>

- Infinite number of DOFs
- Einstein's equations are a system of hyperbolic PDEs

Determinism:

Given all field values within Σ , it is possible to predict uniquely their values anywhere within ${\cal D}$

Classical determinism in General Relativity

Homogeneous cosmologies:

- Infinite \rightarrow finite number of DOFs
- PDEs \rightarrow ODEs

Determinism:

Picard–Lindelöf theorem of existence and uniqueness under a certain set of conditions for the ODEs, an initial-value problem has a unique solution

GR is a gauge Hamiltonian system: **not all degrees of freedom are physical**, and determinism fails only if there is no way to evolve uniquely all physical DOFs


The Bianchi-IX universe

The BIX universe is a spatially **homogeneous** but not necessairly isotropic universe with a 3-sphere topology

$$\mathcal{H} = \frac{1}{2} \left(-k_0^2 + k_1^2 + k_2^2 \right) + \frac{1}{2} e^{\frac{2x^0}{\sqrt{3}}} U(x^1, x^2)$$

- * x^0 scale variable (function of volume v) Big Bang singularity $x^0 \to -\infty$ ($v \to 0$) reached in a finite amount of proper time
- ◆ x^1, x^2 shape variables (measure large-scale anisotropy) when $x^1 = x^2 = 0 \rightarrow$ round 3-sphere metric

The Shape potential




kinetic energy \simeq potential energy \rightarrow **bouces against the potential walls** (Taub transitions) kinetic energy \gg potential energy \rightarrow **straight uniform motion** (Kasner epoch)

Cosmological models evolving through the Big Bang | Martina Adamo, 9 September 2022

41.00

Essential singularity at the BB

Infinite bounces in a finite amount of proper time \rightarrow eternal chaotic behaviour



Observable variable $\theta = \arctan \frac{x^2}{x^1}$ oscillates infinitely fast near the BB BB of the BIX universe is an essential singularity (as $\lim_{x\to 0} \sin \frac{1}{x}$)

Cosmological models evolving through the Big Bang | Martina Adamo, 9 September 2022

Quiescence

Adding an **homogeneous scalar field** whose potential does not grow too fast


Compatible with inflation;

$$\mathcal{H} = \frac{1}{2} \left(-k_0^2 + k_1^2 + k_2^2 + \pi_{\phi}^2 \right) + \frac{1}{2} e^{\frac{2x^0}{\sqrt{3}}} U(x^1, x^2) + e^{\sqrt{3}x^0} V(\phi) \right)$$


After a finite number of bounces, the **bounces will stop**, and the solution will settle on a **single straight-line (Kasner) solution**

0 H C

Shape-plane topology




GR is blind under changes of space orientation \rightarrow two shape planes with different orientations




4.2.20

GR is blind under changes of space orientation \rightarrow two shape planes with different orientations




4110



- Considering two shape planes with opposite orientations
- Compactifing phase-spaces via gnomonic projection on a 2-sphere
- Each shape plane is projected into an hemisphere

 $0 \le \beta < \frac{\pi}{2}, \quad \frac{\pi}{2} < \beta \le \pi$

- Kasner solutions are projected into great circles on the 2-sphere
- BB singularity $\beta \to \pm \frac{\pi}{2}$



Continuation through the BB

Compactifing all physical DOFs y_i , the EOMs satisfy the Picard-Lindelöf Theorem

$$\frac{dy_i}{d\beta} = f_i(y)$$

 $f_i(y)$ are differentiable functions (a stronger property than the Lipschitz-continuity required by the theorem)

Each BIX solution on one side of the singularity is associated with **one and only one** BIX solution on the other side of the singularity

Is the orientation change physical? To detect an orientation change we need matter fields

GR minimally coupled with:

- U(1)-gauge fields
- SU(2)-gauge fields
- a 1-component SU(3)-gauge field

- Quiescence is preserved
- EOMs are still well-behaved at the singularity
 - Gauge-fields do not flip their orientation

EOMs, describing gravity minimally coupled with stiff-matter and with YM fields, satify a theorem of existence and uniqueness at the BB singularity of a BIX universe: each solution passes through the singularity without loss of informations

YM-fields detect the orientation change

- Generalization to a generic SU(3)-YM field
- Adding fermionic fields
- Including inhomogeneities as perturbations about homogeneous terms

MA, F. Mercati, "Continuation through a Big Bang Singularity of the Einstein–Maxwell/Yang–Mills theory", to appear (2022)

F. Mercati, D. Sloan, "Through a Black Hole Singularity", arXiv:2109.10753 [gr-qc] (2021)

F. Mercati, "Through the Big Bang in inflationary cosmology", JCAP 10 025, arXiv:1906.08835 [gr-qc] (2019)


D. Sloan, "Scalar fields and the FLRW singularity", Class. Quant. Grav. 36, 23, 235004, arXiv:1907.08287 (2019)

F. Mercati, "Shape Dynamics: Relativity and Relationalism", Oxford University Press, arXiv:1409.0105 [gr-qc] (2019)

T. Koslowski, F. Mercati, D. Sloan, "Relationalism Evolves the Universe Through the Big Bang", *Phys. Lett. B778, 339, arXiv:1607.02460 [gr-qc] (2018)*

Cosmological models evolving through the Big Bang | Martina Adamo, 9 September 2022

Thank you for the attention!



NASA, ESA and the Hubble Heritage Team (STScI/AURA)