


# Conserved charges from dynamics in non commutative spaces

DOMENICO FRATTULLILLO  
UNIVERSITÀ DI NAPOLI  
FEDERICO II

COST CA18108 Workshop  
on theoretical aspects of  
quantum gravity – Belgrade

In collaboration with:

G. Amelino-Camelia, G. Fabiano



# Non commutative space-times

---

In many quantum gravity models non commutative space-times are considered where an inverse energy scale is introduced  $\ell \simeq l_p$ .

In the last years people have mainly studied the kinematics inspired from the non-commutative framework.

We will investigate particle dynamics on these non-commutative spaces, gaining some insights on the conserved charges for multiparticle systems.

# Hopf-Algebra Symmetries for empty 1+1D $\kappa$ -Minkowski

Coproducts (bicross-product basis)

$$\Delta P_0 = P_0 \otimes 1 + 1 \otimes P_0 \quad \Delta P_1 = P_1 \otimes 1 + e^{-\ell P_0} \otimes P_1 \quad \Delta N = N \otimes 1 + e^{-\ell P_0} \otimes N$$

Commutators

$$[P_0, P_1] = 0 \quad [N, P_0] = iP_1 \quad [N, P_1] = \frac{i}{2\ell} (1 - e^{-2\ell P_0}) - \frac{i\ell}{2} P_1^2$$

Casimir Operator

$$C = \frac{4}{\ell^2} \sinh^2 \left( \frac{\ell}{2} P_0 \right) - (P_1)^2 e^{\ell P_0}$$

Non-commutative space-time

$$[x_1, x_0] = i\ell x_1$$

# Total momentum

---

The standard interpretation is to regard the generators  $P_0, P_1, N$  as conserved charges of free particle motion.

If more than one particle is considered the coproducts usually inspire the composition law of said charges from which we obtain the total momentum:

$$(p \oplus k)_0 = p_0 + k_0$$

$$(p \oplus k)_1 = p_1 + k_1 + e^{-\ell p_0} k_1$$

$$(N_p \oplus N_k) = N_p + N_k + e^{-\ell p_0} N_k$$

# Ambiguities for the total momentum

---

However, this framework suffers from ambiguities in defining such total momentum.

A first ambiguity comes from the fact that in the free particle case, even just the single momentum of each particle is a conserved charge.

And even if we construct the total momentum using the  $\oplus$  inspired by the coproducts there are ambiguities due to the non commutativity of the composition law:

$$(p \oplus k)_1 \neq (k \oplus p)_1$$

What can dynamics tell us about total momentum?

# Spatial “(2+0)D” $\kappa$ -Minkowski (Vanilla) (1)

Let us consider the following map:

$$\ell \rightarrow i\ell \quad x_0 \rightarrow ix_2 \quad P_0 \rightarrow -iP_2 \quad N \rightarrow -iR$$

The Hopf Algebra becomes:

$$[P_1, P_2] = 0 \quad [R, P_1] = \frac{i}{2\ell} (1 - e^{-2\ell P_2}) + \frac{i\ell}{2} P_1^2 \quad [R, P_2] = -iP_1$$

$$\Delta P_1 = P_1 \otimes 1 + e^{-\ell P_2} \otimes P_1 \quad \Delta P_2 = P_2 \otimes 1 + 1 \otimes P_2$$

$$\Delta R = R \otimes 1 + e^{-\ell P_2} \otimes R$$

# Spatial “(2+0)D” $\kappa$ -Minkowski (2)

---

The Casimir operator now reads

$$C = \frac{4}{\ell^2} \sinh^2 \left( \frac{\ell}{2} P_2 \right) + (P_1)^2 e^{\ell P_2} \xrightarrow{\ell \rightarrow 0} P_1^2 + P_2^2$$

and the coordinate non-commutativity becomes

$$[x_1, x_2] = i\ell x_2$$

Now  $x_0$  is a commutative variable and we can set up standard Hamiltonian analysis in the deformed Galilean Relativistic regime.

Furthermore, to ensure Jacobi identities are satisfied, the symplectic structure is also deformed

$$[P_1, x_1] = i \quad [P_2, x_2] = i \quad [P_1, x_2] = 0 \quad [P_2, x_1] = -i\ell P_1$$

# The elastic potential

---

In standard QM, a system of two particles interacting via an elastic potential is described by the following Hamiltonian:

$$H = \frac{\vec{p}_A^2}{2m} + \frac{\vec{p}_B^2}{2m} + \frac{1}{2}\rho(\vec{q}_A - \vec{q}_B)^2$$

The conserved charges read

$$\vec{P} = \vec{p}_A + \vec{p}_B$$

$$\vec{L} = \vec{L}_A + \vec{L}_B$$

$$\vec{L}_I = \vec{x}_I \wedge \vec{p}_I$$

Indeed

$$[H, \vec{L}] = [H, \vec{P}] = 0$$

Our objective will be to construct a deformed Hamiltonian that commutes with total momentum and total angular momentum, as inspired from non-commutativity.



# The deformed Hamiltonian

Keeping our analysis up to second order in the deformation parameter  $\ell^2$ , the deformed kinetic term is given by

$$H_K^I = \frac{(p_1^I)^2}{2m} + \frac{(p_2^I)^2}{2m} + \ell \frac{(p_1^I)^2 p_2^I}{2m} + \frac{\ell^2 (p_1^I)^2 (p_2^I)^2}{4m} + \frac{\ell^2 (p_2^I)^4}{24m}$$

while the most general ansatz for the quadratic potential reads

$$V(\vec{x}^A, \vec{x}^B) = \frac{1}{2} \rho (\vec{x}^A - \vec{x}^B)^2 + \ell \rho \sum \alpha_{ijk}^{IJK} p_i^I x_j^J x_k^K + \ell^2 \rho \sum \beta_{ijkh}^{IJKH} p_i^I p_j^J x_k^K x_h^H$$

The full Hamiltonian for two particles is simply  $H = H_K^A + H_K^B + V(\vec{x}^A, \vec{x}^B)$

# Vanilla $\kappa$ -Minkowski : two particle dynamics

Adopting the usual physical interpretation, for the total charges to be symmetries of the dynamics we require

$$[(p^A \oplus p^B)_1, H] = 0 \quad [(p^A \oplus p^B)_2, H] = 0 \quad [(R^A \oplus R^B), H] = 0$$

This is equivalent to a system of equations involving coefficients  $\alpha$  and  $\beta$ . At first order in  $\ell$ , we find the following solution

$$H = H_K^A + H_K^B + \frac{1}{2}\rho(\vec{x}_A - \vec{x}_B)^2 + \frac{1}{2}\ell\rho[p_2^A x_1^A(x_1^A - x_1^B) - (p_1^A x_1^A - 2p_1^B x_1^B)(x_2^A - x_2^B)] + O(\ell^2)$$

# Vanilla $\kappa$ -Minkowski : three particle dynamics

A possible separable three particle Hamiltonian for this model simply reads

$$H^{ABC} = H_K^A + H_K^B + H_K^C + V(\vec{x}^A, \vec{x}^B) + V(\vec{x}^A, \vec{x}^C) + V(\vec{x}^B, \vec{x}^C) + O(\ell^2)$$

And it can be explicitly checked that

$$[(p^I \oplus p^J \oplus p^K)_i, H^{ABC}] \neq 0 \quad [(R^I \oplus R^J \oplus R^K)_i, H^{ABC}] \neq 0 \quad I, J, K = A, B, C \quad i = 1, 2$$

Even the most general ansatz for a non-separable Hamiltonian fails  $\Rightarrow$  No three-particle dynamics in spatial  $\kappa$ -Minkowski space with coproduct inspired composition laws!

# Proper-dS framework

Momentum space model with  $\kappa$ -Minkowski algebra structure but composition laws given by

$$(p^A \oplus p^B)_1 = p_1^A + p_1^B - \ell(p_2^A p_1^B + p_1^A p_2^B) + \frac{\ell^2}{2} [(p_2^A p_1^B + p_1^A p_2^B)(p_2^A + p_2^B) - p_1^A (p_1^B)^2 - (p_1^A)^2 p_1^B]$$

$$(p^A \oplus p^B)_2 = p_2^A + p_2^B + \ell p_1^A p_1^B - \frac{\ell^2}{2} [(p_1^A)^2 p_2^B + (p_1^B)^2 p_2^A - p_1^A p_1^B (p_2^A + p_2^B)]$$

$$(R^A \oplus R^B) = R^A + R^B$$

The composition law for momentum is commutative but non-associative.

# Proper-dS: dynamics

Requiring that the total charges are symmetries of the dynamics, we obtain infinitely many solutions up to second order in  $\ell^2 \Rightarrow$  A potential  $V(\vec{x}^A, \vec{x}^B)$  exists.

The three-particle separable Hamiltonian

$$H^{ABC} = H_K^A + H_K^B + H_K^C + V(\vec{x}^A, \vec{x}^B) + V(\vec{x}^A, \vec{x}^C) + V(\vec{x}^B, \vec{x}^C)$$

doesn't work with total charges  $((p^A \oplus p^B) \oplus p^C)_i$  nor  $(p^A \oplus (p^B \oplus p^C))_i$ . If we add a non-separable term of the form

$$H^{NS} = \frac{1}{2} l^2 \rho [(p_2^C p_1^A - p_1^C p_2^A)(x_2^C x_1^B - x_1^C x_2^B) + p_2^B (p_1^C x_1^C x_2^A - p_1^C x_2^C x_1^A + p_1^A (p_2^A p_1^C (2x_1^C - x_1^A - x_1^B) - 2x_1^C x_2^C + x_1^C x_2^B + x_2^C x_1^A) + p_1^B (p_2^C (x_2^C x_1^A - x_1^C x_2^A) + p_1^A x_2^C (2x_2^C - x_2^A - x_2^B) + p_2^A x_1^C (x_2^A - 2x_2^C) + p_2^A x_2^C x_1^B]$$

$((p^A \oplus p^B) \oplus p^C)_i$  is conserved.

# Conclusions

---

With coproduct inspired composition law, we can not find a compatible Hamiltonian for the three particle system.

If we consider the proper dS scenario we can construct a non separable Hamiltonian compatible with the three particle system.

To conclude we can say that the Hamiltonian is the fundamental element in order to define the total momentum of a multiparticle system.

**Thank you!**